

TECHNICAL NOTES

Wagg & Purr® HCA Spray



Active Constituents

Hydrocortisone Aceponate 0.584 mg/mL

Actions

Wagg & Purr® HCA Spray is for the treatment of inflammatory and pruritic skin conditions.

Indications

Wagg & Purr® HCA Spray is for the symptomatic relief of inflammatory and pruritic skin conditions. It aids in the reduction of dermatological signs in localised lesions associated with flea allergy dermatitis.

Restrains

DO NOT USE on food producing animals.

Contraindications

This product is contraindicated for use on cutaneous ulcers.

Precautions

In case of concurrent lesions of bacterial, fungal or parasitic origin, the infectious condition should be controlled by an adequate treatment, the use of which (time, duration, dosage) being under prescription of the veterinarian.

In the absence of specific information, the use in animals suffering from Cushing's syndrome shall be based on the risk-benefit assessment.

Since glucocorticoids are known to slow growth, use in young animals (under 7 months of age) shall be based on the risk-benefit assessment and subject to regular clinical evaluations.

Care should be taken to avoid spraying into the eyes of the animal.

The safety of the product has not been evaluated in breeding, pregnant or lactating dogs and it is therefore not recommended.

This product has not been evaluated for extended use. For conditions requiring more than 7 days consecutive use, the responsible veterinarian should subject the use of the product to a risk-benefit assessment taking into consideration the effects of long-term corticosteroid administration.

Total body surface treated should not exceed a surface corresponding to the treatment of two flanks from spine to mammary chains including the shoulders and thighs (1/3rd of the dogs' body).

Dosage & Administration

Wagg & Purr® HCA Spray is for topical administration only. There is no need to rub in.

Apply the product by activating the pump spray from about 10 cm above the affected area.

The recommended dose is 1.52 µg of hydrocortisone aceponate/cm² of affected skin per day. This dosage is achieved with two pump spray activations over a surface area to be treated equivalent to a 10 cm x 10 cm².

Where lesion is less than 10 cm x 10 cm², the prescribing veterinarian may consider adjusting the dose down.

Repeat the treatment once daily for 7 consecutive days.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre on 13 1126. May cause allergy. Warning - May cause birth defects.

Safety Data Sheet

For Safety Data Sheet see www.avet.health

Safety Information

Will irritate eyes. Avoid contact with eyes. If product in eyes, wash out immediately with water. Wash hands after use.

Spray in a well-ventilated area. Flammable. Do not spray on naked flame or incandescent material or smoke while handling.

Presentation

76 mL spray bottle

Disposal

Dispose of empty container by wrapping with paper and putting in garbage.

Storage

Store below 30 °C (room temperature). Protect from light.

Poisons schedule

S4

Registration Numbers

APVMA Approval Number: 91329

Wagg&Purr® is a registered trademark of AVet Health Ltd